

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

7-17-2009

Scout for Corn Leaf Aphids Before Tasseling

Erin W. Hodgson

Iowa State University, ewh@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Hodgson, Erin W., "Scout for Corn Leaf Aphids Before Tasseling" (2009). *Integrated Crop Management News*. Paper 594.
<http://lib.dr.iastate.edu/cropnews/594>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

[Subscribe to Crop News](#)

Archives

[2015](#)[2014](#)[2013](#)[2012](#)[2011](#)[2010](#)[2009](#)[2008](#)[Previous Years](#)

ISU Crop Resources

[Extension Field Agronomists](#)[Crop & Soils Info](#)[Pesticide Applicator Training](#)[Agronomy Extension](#)[Entomology Extension](#)[Plant Pathology Extension](#)[Ag and Biosystems Engineering Extension](#)[Agribusiness Education Program](#)[Iowa Grain Quality Initiative](#)[College of Agriculture and Life Sciences](#)[ISU Extension](#)

Integrated Crop Management NEWS

[PRINT STORY](#)
[EMAIL STORY](#)
[ADD TO DELICIOUS](#)
[ATOM FEED](#)
[FOLLOW ON TWITTER](#)

Scout for Corn Leaf Aphids Before Tasseling

By Erin Hodgson, Department of Entomology

Corn leaf aphids can be common in Iowa corn, but they rarely cause significant economic losses. They prefer sorghum as a host plant, but will also feed on corn, barley, millet and many grasses. Although infestations become apparent when tassels emerge, colonies can start to develop in late June.

Description and life cycle. Corn leaf aphids are bluish-green in color with dark legs and cornicles (tailpipes at the end of the abdomen), and usually less than one-eighth inch in length. Like all aphids, they have a piercing-sucking mouthpart that feeds on plant phloem. Corn leaf aphids excrete a sugar-rich honeydew that can cover corn leaves, tassels and silks.



Corn leaf aphids and cast skins from molting.

The corn leaf aphid is not known to overwinter in Iowa, but migrates on airstreams each season. Females give live birth to nymphs via asexual reproduction during the summer. Wingless forms are more common, however, migratory winged forms are produced during crowded conditions.



Heavy corn leaf aphid infestation.

Damage. Feeding by large aphid colonies can reduce yields significantly. Aphids feed on upper leaves, in the whorl or on unemerged tassels. Heavy corn leaf aphid infestations may cause leaves to curl, wilt and become yellow. Tassels and silks can become covered with honeydew. Aphid colonies can slowly kill the tassel and top leaves of a corn plant. The plant responds by shunting nutrients to the feeding area and not to the developing ear, resulting in stunted ears or even barren plants. The most severe damage occurs between the late-whorl and pollination stages.

Scouting and management. Scouting for corn leaf aphids must be done before tasseling - ideally three weeks before. Developing colonies are most commonly found deep within the whorl. Stop at five locations and examine twenty plants. Carefully pull the whorl leaves away from the stalk and unroll the leaves to estimate the number of aphids. Closer to tasseling, aphids move from within the whorl and are very exposed to weather and natural enemies. Take note of beneficial insects (e.g., ladybeetles and lacewings) and/or parasitized mummies; consider not treating aphids if more than 20 percent are parasitized.

Spraying for corn leaf aphids while they are protected deep in the whorl is not effective. Treating for corn leaf aphids after tasseling is usually not a cost-effective management decision because aphid populations will decline naturally. A well-timed spray while aphids are outside the whorl or on the tassel is recommended when they exceed thresholds. Corn fields can be particularly sensitive to corn leaf aphid feeding if they are drought-stressed. If 50 percent of the plants have more than 100 aphids per plant, tassels are coated in honeydew and plants are under drought stress, treatment may be justified. There are several products registered in Iowa for corn leaf aphid (Table 1). Follow label directions and pay attention to spray guidelines.

Table 1. Insecticides labeled for corn leaf aphid in corn

<i>Product (active ingredient)</i>	<i>Application rate</i>	<i>Preharvest interval</i>
Adjourn (esfenvalerate)	5.8-9.6 oz/ac	21
Proaxis (gamma-cyhalothrin)	2.56-3.84 oz/ac	21
Thionex 3EC (endosulfan)	1.33 qts/ac	1 (sweet corn only)
Warrior II (lambda-cyhalothrin)	1.28-1.92 oz/ac	21

Portions of this article were originally appeared here:

<http://www.ipm.iastate.edu/ipm/icm/1998/8-10-1998/cornaphids.html>

Erin Hodgson is an assistant professor of entomology with extension and research responsibilities. She can be contacted by email at ewh@iastate.edu or phone (515) 294-2847.

This article was published originally on 7/17/2009. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.